

REFINE

This result exported from <http://refinenutrition.org> on July 20, 2017 at 8:29 PM

Filters Applied

Efficacy of iron-supplement bars to reduce anemia in urban Indian women: a cluster-randomized controlled trial

URL: <http://refinenutrition.org/content/efficacy-iron-supplement-bars-reduce-anemia-urban-indian-women-cluster-randomized-controlled>

Improved nutrition in the first 1000 days and adult human capital and health

URL: <http://refinenutrition.org/content/improved-nutrition-first-1000-days-and-adult-human-capital-and-health>

A 2-year integrated agriculture and nutrition and health behavior change communication program targeted to women in Burkina Faso reduces anemia, wasting, and diarrhea in children 3-12.9 months of age at baseline: a cluster-randomized controlled trial

URL: <http://refinenutrition.org/content/2-year-integrated-agriculture-and-nutrition-and-health-behavior-change-communication-program>

A cluster RCT to measure the effectiveness and of cash-based interventions on nutrition status, Sindh Province, Pakistan

URL: <http://refinenutrition.org/content/cluster-rct-measure-effectiveness-and-cash-based-interventions-nutrition-status-sindh>

A cluster-randomized trial determining the efficacy of caterpillar cereal as a locally available and sustainable complementary food to prevent stunting and anaemia

URL: <http://refinenutrition.org/content/cluster-randomized-trial-determining-efficacy-caterpillar-cereal-locally-available-and>

A combined intervention of zinc, multiple micronutrients, and Albendazole does not ameliorate environmental enteric dysfunction or stunting in rural Malawian children in a double-blind randomized controlled trial

URL: <http://refinenutrition.org/content/combined-intervention-zinc-multiple-micronutrients-and-albendazole-does-not-ameliorate>

A comprehensive linear programming tool to optimize formulations of ready-to-use therapeutic foods: an application to Ethiopia

URL: <http://refinenutrition.org/content/comprehensive-linear-programming-tool-optimize-formulations-ready-use-therapeutic-foods>

A food transfer program without a formal education component modifies complementary feeding practices in poor rural Mexican communities

URL: <http://refinenutrition.org/content/food-transfer-program-without-formal-education-component-modifies-complementary-feeding>

A lipid-based nutrient supplement but not corn-soy blend modestly increases weight gain among 6-to 18-month-old moderately underweight children in rural Malawi

URL: <http://refinenutrition.org/content/lipid-based-nutrient-supplement-not-corn-soy-blend-modestly-increases-weight-gain-among-6-18>

A mixed method study exploring adherence to and acceptability of small quantity lipid-based nutrient supplements (SQ-LNS) among pregnant and lactating women in Ghana and Malawi

URL: <http://refinenutrition.org/content/mixed-method-study-exploring-adherence-and-acceptability-small-quantity-lipid-based-nutrient>

A novel fortified blended flour, corn-soy blend "plus-plus," is not inferior to lipid-based ready-to-use supplementary foods for the treatment of moderate acute malnutrition in Malawian children

URL: <http://refinenutrition.org/content/novel-fortified-blended-flour-corn-soy-blend-plus-plus-not-inferior-lipid-based-ready-use>

A randomized controlled facility based trial to assess the impact of indigenously prepared ready to use therapeutic food (RUTF) for children with severe acute malnutrition in India

URL: <http://refinenutrition.org/content/randomized-controlled-facility-based-trial-assess-impact-indigenously-prepared-ready-use>

A randomized controlled trial offering higher- compared with lower-dairy second meals daily in preschools in Guinea-Bissau demonstrates an attendance-dependent increase in weight gain for both meal types and an increase in mid-upper arm circumference

URL: <http://refinenutrition.org/content/randomized-controlled-trial-offering-higher-compared-lower-dairy-second-meals-daily>

A ready-to-use therapeutic food containing 10% milk is less effective than one with 25% milk in the treatment of severely malnourished children

URL: <http://refinenutrition.org/content/ready-use-therapeutic-food-containing-10-milk-less-effective-one-25-milk-treatment-severely>

A single-centre, randomized, single-blind, parallel group clinical trial in rural Malawi, testing the growth promoting effect of long-term complementary feeding of infants with a high-energy, micronutrient fortified spread

URL: <http://refinenutrition.org/content/single-centre-randomized-single-blind-parallel-group-clinical-trial-rural-malawi-testing>

A tale of two measures: Self-report and lab-assessed values in amount of oil Added to CSB porridge prepared by caregivers of children with moderate acute malnutrition in Southern Malawi

URL: <http://refinenutrition.org/content/tale-two-measures-self-report-and-lab-assessed-values-amount-oil-added-csb-porridge-prepared>

Acceptability and efficacy of locally produced ready-to-use therapeutic food nutreal in the management of severe acute malnutrition in comparison with defined food: A randomized control trial

URL: <http://refinenutrition.org/content/acceptability-and-efficacy-locally-produced-ready-use-therapeutic-food-nutreal-management>

Acceptability of a lipid-based nutrient supplement among Guatemalan infants and young children

URL: <http://refinenutrition.org/content/acceptability-lipid-based-nutrient-supplement-among-guatemalan-infants-and-young-children>

Acceptability of lipid-based nutrient supplements (LNS) among Ghanaian infants and pregnant or lactating women

URL: <http://refinenutrition.org/content/acceptability-lipid-based-nutrient-supplements-lns-among-ghanaian-infants-and-pregnant-or>

Acceptability of lipid-based nutrient supplements and micronutrient powders among pregnant and lactating women and infants and young children in bangladesh and their perceptions about malnutrition and nutrient supplements

URL: <http://refinenutrition.org/content/acceptability-lipid-based-nutrient-supplements-and-micronutrient-powders-among-pregnant-and>

Acceptability of locally produced ready-to-use therapeutic foods in Ethiopia, Ghana, Pakistan and India

URL: <http://refinenutrition.org/content/acceptability-locally-produced-ready-use-therapeutic-foods-ethiopia-ghana-pakistan-and-india>

Acceptability of new formulations of corn-soy blends and lipid-based nutrient supplements in Province du Passor, Burkina Faso

URL: <http://refinenutrition.org/content/acceptability-new-formulations-corn-soy-blends-and-lipid-based-nutrient-supplements-province>

Acceptability of novel small-quantity lipid-based nutrient supplements for complementary feeding in a peri-urban South African community

URL: <http://refinenutrition.org/content/acceptability-novel-small-quantity-lipid-based-nutrient-supplements-complementary-feeding>

Acceptability of outpatient ready-to-use food-based protocols in HIV-infected Senegalese children and adolescents within the MAGGEN cohort study

URL: <http://refinenutrition.org/content/acceptability-outpatient-ready-use-food-based-protocols-hiv-infected-senegalese-children-and>

Acceptability of three novel lipid-based nutrient supplements among Malawian infants and their caregivers

URL: <http://refinenutrition.org/content/acceptability-three-novel-lipid-based-nutrient-supplements-among-malawian-infants-and-their>

Acceptability of zinc-fortified, lipid-based nutrient supplements (LNS) prepared for young children in Burkina Faso

URL: <http://refinenutrition.org/content/acceptability-zinc-fortified-lipid-based-nutrient-supplements-lns-prepared-young-children>

Adding a water, sanitation and hygiene intervention and a lipid-based nutrient supplement to an integrated agriculture and nutrition program improved the nutritional status of young Burkinab children

URL: <http://refinenutrition.org/content/adding-water-sanitation-and-hygiene-intervention-and-lipid-based-nutrient-supplement>

Adding multiple micronutrient powders to a homestead food production programme yields marginally significant benefit on anaemia reduction among young children in Nepal

URL: <http://refinenutrition.org/content/adding-multiple-micronutrient-powders-homestead-food-production-programme-yields-marginally>

Age-based preventive targeting of food assistance and behaviour change and communication for reduction of childhood undernutrition in Haiti: a cluster randomised trial

URL: <http://refinenutrition.org/content/age-based-preventive-targeting-food-assistance-and-behaviour-change-and-communication>

An effectiveness trial showed lipid-based nutrient supplementation but not corn-soya blend offered a modest benefit in weight gain among 6- to 18-month-old underweight children in rural Malawi

URL: <http://refinenutrition.org/content/effectiveness-trial-showed-lipid-based-nutrient-supplementation-not-corn-soya-blend-offered>

Antenatal dietary education and supplementation to increase energy and protein intake

URL: <http://refinenutrition.org/content/antenatal-dietary-education-and-supplementation-increase-energy-and-protein-intake>

Antibiotics as part of the management of severe acute malnutrition

URL: <http://refinenutrition.org/content/antibiotics-part-management-severe-acute-malnutrition>

Assessment of outpatient therapeutic programme for severe acute malnutrition in three regions of Ethiopia

URL: <http://refinenutrition.org/content/assessment-outpatient-therapeutic-programme-severe-acute-malnutrition-three-regions-ethiopia>

Assessment of the effectiveness of a small quantity lipid-based nutrient supplement on reducing anaemia and stunting in refugee populations in the Horn of Africa: Secondary data analysis

URL: <http://refinenutrition.org/content/assessment-effectiveness-small-quantity-lipid-based-nutrient-supplement-reducing-anaemia-and>

Balancing omega-6 and omega-3 fatty acids in ready-to-use therapeutic foods (RUTF)

URL: <http://refinenutrition.org/content/balancing-omega-6-and-omega-3-fatty-acids-ready-use-therapeutic-foods-rutf>

Cash, food, or vouchers? Evidence from a randomized experiment in northern Ecuador

URL: <http://refinenutrition.org/content/cash-food-or-vouchers-evidence-randomized-experiment-northern-ecuador>

Catch-up growth in children with severe acute malnutrition: comparison of two therapeutic feeding approaches at a district hospital, South Africa

URL: <http://refinenutrition.org/content/catch-growth-children-severe-acute-malnutrition-comparison-two-therapeutic-feeding>

Caterpillar cereal as a potential complementary feeding product for infants and young children: nutritional content and acceptability

URL: <http://refinenutrition.org/content/caterpillar-cereal-potential-complementary-feeding-product-infants-and-young-children>

Cereals and pulse-based ready-to-use therapeutic food as an alternative to the standard milk- and peanut paste-based formulation for treating severe acute malnutrition: a noninferiority, individually randomized controlled efficacy clinical trial

URL: <http://refinenutrition.org/content/cereals-and-pulse-based-ready-use-therapeutic-food-alternative-standard-milk-and-peanut>

Challenges in implementing the integrated community-based outpatient therapeutic program for severely malnourished children in rural Southern Ethiopia

URL: <http://refinenutrition.org/content/challenges-implementing-integrated-community-based-outpatient-therapeutic-program-severely>

Changes in plasma phosphate during in-patient treatment of children with severe acute malnutrition: an observational study in Uganda

URL: <http://refinenutrition.org/content/changes-plasma-phosphate-during-patient-treatment-children-severe-acute-malnutrition>

Changes in whole-blood PUFA and their predictors during recovery from severe acute malnutrition

URL: <http://refinenutrition.org/content/changes-whole-blood-pufa-and-their-predictors-during-recovery-severe-acute-malnutrition>

Children successfully treated for moderate acute malnutrition remain at risk for malnutrition and death in the subsequent year after recovery

URL: <http://refinenutrition.org/content/children-successfully-treated-moderate-acute-malnutrition-remain-risk-malnutrition-and-death>

Clinical and anthropometric profile of severe acute malnourished children on therapeutic intervention with WHO/UNICEF recommended therapeutic food

URL: <http://refinenutrition.org/content/clinical-and-anthropometric-profile-severe-acute-malnourished-children-therapeutic>

Clinical significance of neuropsychological improvement after supplementation with omega-3 in 812 years old malnourished Mexican children: A randomized, double-blind, placebo and treatment clinical trial

URL: <http://refinenutrition.org/content/clinical-significance-neuropsychological-improvement-after-supplementation-omega-3-812-0>

Cofortification of ferric pyrophosphate and citric acid/trisodium citrate into extruded rice grains doubles iron bioavailability through in situ generation of soluble ferric pyrophosphate citrate complexes

URL: <http://refinenutrition.org/content/cofortification-ferric-pyrophosphate-and-citric-acidtrisodium-citrate-extruded-rice-grains>

Combined protocol for SAM/MAM treatment: The ComPAS study

URL: <http://refinenutrition.org/content/combined-protocol-sammam-treatment-compas-study>

Community management of acute malnutrition in the developing world

URL: <http://refinenutrition.org/content/community-management-acute-malnutrition-developing-world>

Community-based management of child malnutrition in Zambia: HIV/AIDS infection and other risk factors on child survival

URL: <http://refinenutrition.org/content/community-based-management-child-malnutrition-zambia-hiv-aids-infection-and-other-risk>

Community-based management of severe acute malnutrition in India: new evidence from Bihar

URL: <http://refinenutrition.org/content/community-based-management-severe-acute-malnutrition-india-new-evidence-bihar>

Community-based management of severe malnutrition: SAM and SUW in children under five in the Melghat tribal area, central India

URL: <http://refinenutrition.org/content/community-based-management-severe-malnutrition-sam-and-suw-children-under-five-melghat>

Community-based supplementary feeding for promoting the growth of children under five years of age in low and middle income countries

URL: <http://refinenutrition.org/content/community-based-supplementary-feeding-promoting-growth-children-under-five-years-age-low-and>

Comparative weight gain with ready-to-use therapeutic food in stunted HIV-infected and-uninfected children in a Nigerian Hospital

URL: <http://refinenutrition.org/content/comparative-weight-gain-ready-use-therapeutic-food-stunted-hiv-infected-and-uninfected>

Comparing milk-fortified corn-soy blend (CSB++), soy ready-to-use supplementary food (RUSF), and soy/whey RUSF (Supplementary Plumpy) in the treatment of moderate acute malnutrition

URL: <http://refinenutrition.org/content/comparing-milk-fortified-corn-soy-blend-csb-soy-ready-use-supplementary-food-rusf-and>

Comparison of four different supplementary foods in the treatment of moderate acute malnutrition (MAM) in children under five in Sierra Leone

URL: <http://refinenutrition.org/content/comparison-four-different-supplementary-foods-treatment-moderate-acute-malnutrition-mam>

Comparison of four supplementary foods in treating moderate acute malnutrition in Sierra Leone: an Ebola-constrained cluster-randomized, controlled clinic-based effectiveness trial: A Report from the Food Aid Quality Review

URL: <http://refinenutrition.org/content/comparison-four-supplementary-foods-treating-moderate-acute-malnutrition-sierra-leone-ebola>

Comparison of methods to assess adherence to small-quantity lipid-based nutrient supplements (SQ-LNS) and dispersible tablets among young Burkinab children participating in a community-based intervention trial

URL: <http://refinenutrition.org/content/comparison-methods-assess-adherence-small-quantity-lipid-based-nutrient-supplements-sq-lns>

Comparison of the effectiveness of a milk-free soy-maize-sorghum-based ready-to-use therapeutic food to standard ready-to-use therapeutic food with 25% milk in nutrition management of severely acutely malnourished Zambian children

URL: <http://refinenutrition.org/content/comparison-effectiveness-milk-free-soy-maize-sorghum-based-ready-use-therapeutic-food>

Comparison of the efficacy of a solid ready-to-use food and a liquid, milk-based diet for the rehabilitation of severely malnourished children: A randomized trial

URL: <http://refinenutrition.org/content/comparison-efficacy-solid-ready-use-food-and-liquid-milk-based-diet-rehabilitation-severely>

Comparison of weight-for-height and mid-upper arm circumference (MUAC) in a therapeutic feeding programme in South Sudan: is MUAC alone a sufficient criterion for admission of children at high risk of mortality?

URL: <http://refinenutrition.org/content/comparison-weight-height-and-mid-upper-arm-circumference-muac-therapeutic-feeding-programme>

Considerations in developing lipid-based nutrient supplements for prevention of undernutrition: experience from the International Lipid-Based Nutrient Supplements (iLiNS) Project

URL: <http://refinenutrition.org/content/considerations-developing-lipid-based-nutrient-supplements-prevention-undernutrition>

Cost analysis of the treatment of severe acute malnutrition in West Africa

URL: <http://refinenutrition.org/content/cost-analysis-treatment-severe-acute-malnutrition-west-africa>

Cost-effectiveness of the community-based management of severe acute malnutrition by community health workers in southern Bangladesh

URL: <http://refinenutrition.org/content/cost-effectiveness-community-based-management-severe-acute-malnutrition-community-health>

Dairy products and physical stature: a systematic review and meta-analysis of controlled trials

URL: <http://refinenutrition.org/content/dairy-products-and-physical-stature-systematic-review-and-meta-analysis-controlled-trials>

Delivering improved nutrition: Dairy ingredients in food aid products

URL: <http://refinenutrition.org/content/delivering-improved-nutrition-dairy-ingredients-food-aid-products>

Delivery channels to close the nutrient intake gap What to do without data?

URL: <http://refinenutrition.org/content/delivery-channels-close-nutrient-intake-gapwhat-do-without-data>

Developing food supplements for moderately malnourished children: lessons learned from ready-to-use therapeutic foods

URL: <http://refinenutrition.org/content/developing-food-supplements-moderately-malnourished-children-lessons-learned-ready-use>

Development and acceptability of a novel milk-free soybean-maize-sorghum ready-to-use therapeutic food (SMS-RUTF) based on industrial extrusion cooking process

URL: <http://refinenutrition.org/content/development-and-acceptability-novel-milk-free-soybean-maize-sorghum-ready-use-therapeutic>

Development and acceptability testing of ready-to-use supplementary food made from locally available food ingredients in Bangladesh

URL: <http://refinenutrition.org/content/development-and-acceptability-testing-ready-use-supplementary-food-made-locally-available>

Developmental outcomes among 18monthold Malawians after a year of complementary feeding with lipidbased nutrient supplements or cornsoy flour

URL: <http://refinenutrition.org/content/developmental-outcomes-among-18monthold-malawians-after-year-complementary-feeding-lipid>

Donated fortified cereal blends improve the nutrient density of traditional complementary foods in Haiti, but iron and zinc gaps remain for infants

URL: <http://refinenutrition.org/content/donated-fortified-cereal-blends-improve-nutrient-density-traditional-complementary-foods>

Double-blind cluster randomised controlled trial of wheat flour chapatti fortified with micronutrients on the status of vitamin A and iron in school-aged children in rural Bangladesh

URL: <http://refinenutrition.org/content/double-blind-cluster-randomised-controlled-trial-wheat-flour-chapatti-fortified>

Early invitation to food and/or multiple micronutrient supplementation in pregnancy does not affect body composition in offspring at 54 months: follow-up of the MINIMat randomised trial, Bangladesh

URL: <http://refinenutrition.org/content/early-invitation-food-andor-multiple-micronutrient-supplementation-pregnancy-does-not-affect>

Effect of a nutritional supplement on growth in short and lean prepubertal children: a prospective, randomized, double-blind, placebo-controlled study

URL: <http://refinenutrition.org/content/effect-nutritional-supplement-growth-short-and-lean-prepubertal-children-prospective>

Effect of a package of health and nutrition services on sustained recovery in children after moderate acute malnutrition and factors related to sustaining recovery: a cluster-randomized trial

URL: <http://refinenutrition.org/content/effect-package-health-and-nutrition-services-sustained-recovery-children-after-moderate>

Effect of complementary feeding with lipid-based nutrient supplements and corn-soy blend on the incidence of stunting and linear growth among 6- to 18-month-old infants and children in rural Malawi

URL: <http://refinenutrition.org/content/effect-complementary-feeding-lipid-based-nutrient-supplements-and-corn-soy-blend-incidence>

Effect of complementary feeding with lipid-based nutrient supplements and corn-soy blend on the incidence of stunting and linear growth among 6- to 18-month-old infants and children in rural Malawi

URL: <http://refinenutrition.org/content/effect-complementary-feeding-lipid-based-nutrient-supplements-and-corn-soy-blend-incidence>

Effect of fortified complementary food supplementation on child growth in rural Bangladesh: A cluster-randomized trial

URL: <http://refinenutrition.org/content/effect-fortified-complementary-food-supplementation-child-growth-rural-bangladesh-cluster>

Effect of improved complementary foods on lean body mass, essential fatty acids, and gross motor development of Kenyan infants

URL: <http://refinenutrition.org/content/effect-improved-complementary-foods-lean-body-mass-essential-fatty-acids-and-gross-motor>

Effect of mass supplementation with ready-to-use supplementary food during an anticipated nutritional emergency

URL: <http://refinenutrition.org/content/effect-mass-supplementation-ready-use-supplementary-food-during-anticipated-nutritional>

Effect of preventive supplementation with ready-to-use therapeutic food on the nutritional status, mortality, and morbidity of children aged 6 to 60 months in Niger: a cluster randomized trial.

URL: <http://refinenutrition.org/content/effect-preventive-supplementation-ready-use-therapeutic-food-nutritional-status-mortality>

Effect of ready-to-use foods for preventing child undernutrition in Niger: analysis of a prospective intervention study over 15 months of follow-up

URL: <http://refinenutrition.org/content/effect-ready-use-foods-preventing-child-undernutrition-niger-analysis-prospective>

Effect of short-term supplementation with ready-to-use therapeutic food or micronutrients for children after illness for prevention of malnutrition: a randomised controlled trial in Uganda

URL: <http://refinenutrition.org/content/effect-short-term-supplementation-ready-use-therapeutic-food-or-micronutrients-children>

Effect of supplementation with a lipid-based nutrient supplement on the micronutrient status of children aged 6-18 months living in the rural region of Intibuc, Honduras

URL: <http://refinenutrition.org/content/effect-supplementation-lipid-based-nutrient-supplement-micronutrient-status-children-aged-6>

Effectiveness of a nutritional program based upon a ready-to-use therapeutic food in Chad

URL: <http://refinenutrition.org/content/effectiveness-nutritional-program-based-upon-ready-use-therapeutic-food-chad>

Effectiveness of complementary food supplements and dietary counselling on anaemia and stunting in children aged 6-23 months in poor areas of Qinghai Province, China: a controlled interventional study

URL: <http://refinenutrition.org/content/effectiveness-complementary-food-supplements-and-dietary-counselling-anaemia-and-stunting-0>

Effectiveness of complementary food supplements and dietary counselling on anaemia and stunting in children aged 6-23 months in poor areas of Qinghai Province, China: a controlled interventional study

URL: <http://refinenutrition.org/content/effectiveness-complementary-food-supplements-and-dietary-counselling-anaemia-and-stunting>

Effectiveness of indigenous ready-to-use therapeutic food in community-based management of uncomplicated severe acute malnutrition: a randomized controlled trial from India.

URL: <http://refinenutrition.org/content/effectiveness-indigenous-ready-use-therapeutic-food-community-based-management-uncomplicated>

Effectiveness of milk whey protein-based ready-to-use therapeutic food in treatment of severe acute malnutrition in Malawian under-5 children: a randomised, double-blind, controlled non-inferiority clinical trial

URL: <http://refinenutrition.org/content/effectiveness-milk-whey-protein-based-ready-use-therapeutic-food-treatment-severe-acute>

Effectiveness of nutritional supplementation (RUTF and multi micronutrient) in preventing malnutrition and morbidity in children 6-59 months with infection

URL: <http://refinenutrition.org/content/effectiveness-nutritional-supplementation-rutf-and-multi-micronutrient-preventing>

Effectiveness of three commonly used transition phase diets in the inpatient management of children with severe acute malnutrition: a pilot randomized controlled trial in Malawi

URL: <http://refinenutrition.org/content/effectiveness-three-commonly-used-transition-phase-diets-inpatient-management-children>

Effectiveness study for the development of a home fortification programme for young children in Zambia - endline report

URL: <http://refinenutrition.org/content/effectiveness-study-development-home-fortification-programme-young-children-zambia-endline>

Effects of a lipid-based nutrient supplement during pregnancy and lactation on maternal plasma fatty acid status and lipid profile: Results of two randomized controlled trials

URL: <http://refinenutrition.org/content/effects-lipid-based-nutrient-supplement-during-pregnancy-and-lactation-maternal-plasma-fatty>

Effects of an intervention on infant growth and development: evidence for different mechanisms at work

URL: <http://refinenutrition.org/content/effects-intervention-infant-growth-and-development-evidence-different-mechanisms-work>

Effects of animal source food and micronutrient fortification in complementary food products on body composition, iron status, and linear growth: A randomized trial in Cambodia

URL: <http://refinenutrition.org/content/effects-animal-source-food-and-micronutrient-fortification-complementary-food-products-body>

Effects of community-based follow-up care in managing severely underweight children

URL: <http://refinenutrition.org/content/effects-community-based-follow-care-managing-severely-underweight-children>

Effects of different complementary feeding regimens on iron status and enteric microbiota in breastfed infants

URL: <http://refinenutrition.org/content/effects-different-complementary-feeding-regimens-iron-status-and-enteric-microbiota>

Effects of maternal and child lipid-based nutrient supplements on infant development: a randomized trial in Malawi

URL: <http://refinenutrition.org/content/effects-maternal-and-child-lipid-based-nutrient-supplements-infant-development-randomized>

Effects of pre- and post-natal lipid-based nutrient supplements on infant development in a randomized trial in Ghana

URL: <http://refinenutrition.org/content/effects-pre-and-post-natal-lipid-based-nutrient-supplements-infant-development-randomized>

Effects of responsive stimulation and nutrition interventions on children's development and growth at age 4 years in a disadvantaged population in Pakistan: A longitudinal follow-up of a cluster-randomised factorial effectiveness trial

URL: <http://refinenutrition.org/content/effects-responsive-stimulation-and-nutrition-interventions-childrens-development-and-growth>

Effects of small-quantity lipid-based nutrient supplement on hemoglobin and iron status of pregnant Ghanaian women

URL: <http://refinenutrition.org/content/effects-small-quantity-lipid-based-nutrient-supplement-hemoglobin-and-iron-status-pregnant>

Effects of unconditional cash transfers on the outcome of treatment for severe acute malnutrition (SAM): a cluster-randomised trial in the Democratic Republic of the Congo

URL: <http://refinenutrition.org/content/effects-unconditional-cash-transfers-outcome-treatment-severe-acute-malnutrition-sam-cluster>

Efficacy of sorghum peanut blend and corn soy blend plus in the treatment of moderate acute malnutrition in children aged 659 months in Karamoja, Uganda: a cluster randomized trial

URL: <http://refinenutrition.org/content/efficacy-sorghum-peanut-blend-and-corn-soy-blend-plus-treatment-moderate-acute-malnutrition>

Efficacy of three feeding regimens for home-based management of children with uncomplicated severe acute malnutrition: a randomised trial in India

URL: <http://refinenutrition.org/content/efficacy-three-feeding-regimens-home-based-management-children-uncomplicated-severe-acute>

Enhancing processes for introduction, production, quality assurance, and delivery of US Title II food aid products

URL: <http://refinenutrition.org/content/enhancing-processes-introduction-production-quality-assurance-and-delivery-us-title-ii-food>

Establishing desirable fortificant levels for calcium, iron and zinc in foods for infant and young child feeding: examples from three Asian countries

URL: <http://refinenutrition.org/content/establishing-desirable-fortificant-levels-calcium-iron-and-zinc-foods-infant-and-young-child>

Evaluation of the acceptability of improved supplementary foods for the treatment of moderate acute malnutrition in Burkina Faso using a mixed method approach

URL: <http://refinenutrition.org/content/evaluation-acceptability-improved-supplementary-foods-treatment-moderate-acute-malnutrition>

Evidence of the effectiveness of flour fortification programs on iron status and anemia: a systematic review

URL: <http://refinenutrition.org/content/evidence-effectiveness-flour-fortification-programs-iron-status-and-anemia-systematic-review>

Extending supplementary feeding for children younger than 5 years with moderate acute malnutrition leads to lower relapse rates

URL: <http://refinenutrition.org/content/extending-supplementary-feeding-children-younger-5-years-moderate-acute-malnutrition-leads>

Feasibility and acceptability study of preparing corn soy blend with fortified vegetable oil in Malawi: Formative research

URL: <http://refinenutrition.org/content/feasibility-and-acceptability-study-preparing-corn-soy-blend-fortified-vegetable-oil-malawi>

Feeding behaviors during home-based treatment of moderate acute malnutrition using corn-soy blends or lipid-based nutrient supplements

URL: <http://refinenutrition.org/content/feeding-behaviors-during-home-based-treatment-moderate-acute-malnutrition-using-corn-soy>

Feeding patterns and behaviors during home supplementation of underweight Malawian children with lipid-based nutrient supplements or corn-soy blend

URL: <http://refinenutrition.org/content/feeding-patterns-and-behaviors-during-home-supplementation-underweight-malawian-children>

Follow-up of post-discharge growth and mortality after treatment for severe acute malnutrition (FuSAM study): a prospective cohort study

URL: <http://refinenutrition.org/content/follow-post-discharge-growth-and-mortality-after-treatment-severe-acute-malnutrition-fusam>

Follow-up status of children with SAM treated with RUTF in peri-urban and rural Northern Bangladesh

URL: <http://refinenutrition.org/content/follow-status-children-sam-treated-rutf-peri-urban-and-rural-northern-bangladesh>

Food commodity pipeline management in transitional settings: challenges and lessons learned from the first USAID food development program in South Sudan

URL: <http://refinenutrition.org/content/food-commodity-pipeline-management-transitional-settings-challenges-and-lessons-learned>

Food sharing practices in households receiving supplemental foods for the treatment of moderate acute malnutrition in Ethiopian children

URL: <http://refinenutrition.org/content/food-sharing-practices-households-receiving-supplemental-foods-treatment-moderate-acute>

Food, nutrients, and child growth: The role of specific foods and nutrients in child malnutrition and implications for food assistance programs

URL: <http://refinenutrition.org/content/food-nutrients-and-child-growth-role-specific-foods-and-nutrients-child-malnutrition-and>

Formative assessment to design the packaging of a lipid-based nutrient supplement for a home fortification program to improve the nutritional status of young children in the Democratic Republic of Congo (DRC)

URL: <http://refinenutrition.org/content/formative-assessment-design-packaging-lipid-based-nutrient-supplement-home-fortification>

Formative research for the development of a marketbased home fortification programme for young children in Niger

URL: <http://refinenutrition.org/content/formative-research-development-marketbased-home-fortification-programme-young-children>

Formulation of complementary food using amaranth, chickpea and maize improves iron, calcium and zinc content

URL: <http://refinenutrition.org/content/formulation-complementary-food-using-amaranth-chickpea-and-maize-improves-iron-calcium-and>

Fortification of staple foods with zinc for improving zinc status and other health outcomes in the general population.

URL: <http://refinenutrition.org/content/fortification-staple-foods-zinc-improving-zinc-status-and-other-health-outcomes-general>

Fortified snack reduced anemia in rural school-aged children of Haiti: a cluster-randomized, controlled trial

URL: <http://refinenutrition.org/content/fortified-snack-reduced-anemia-rural-school-aged-children-haiti-cluster-randomized>

Gut DNA viromes of Malawian twins discordant for severe acute malnutrition

URL: <http://refinenutrition.org/content/gut-dna-viromes-malawian-twins-discordant-severe-acute-malnutrition>

Gut microbiomes of Malawian twin pairs discordant for kwashiorkor

URL: <http://refinenutrition.org/content/gut-microbiomes-malawian-twin-pairs-discordant-kwashiorkor>

Gut microbiota in Malawian infants in a nutritional supplementation trial

URL: <http://refinenutrition.org/content/gut-microbiota-malawian-infants-nutritional-supplementation-trial>

Gut microbiota of healthy and malnourished children in Bangladesh

URL: <http://refinenutrition.org/content/gut-microbiota-healthy-and-malnourished-children-bangladesh>

High awareness but low coverage of a locally produced fortified complementary food in Abidjan, Cte dlvoire: Findings from a cross-sectional survey

URL: <http://refinenutrition.org/content/high-awareness-low-coverage-locally-produced-fortified-complementary-food-abidjan-cte-d>

High oleic ready-to-use therapeutic food maintains docosahexaenoic acid status in severe malnutrition: A randomized, blinded trial

URL: <http://refinenutrition.org/content/high-oleic-ready-use-therapeutic-food-maintains-docosahexaenoic-acid-status-severe>

Hospital management of severe acute malnutrition in children with F-75 and F-100 alternative local preparations: results and challenges

URL: <http://refinenutrition.org/content/hospital-management-severe-acute-malnutrition-children-f-75-and-f-100-alternative-local>

Identifying the sociocultural barriers and facilitating factors to nutrition-related behavior change formative research for a stunting prevention program in Ntchisi, Malawi

URL: <http://refinenutrition.org/content/identifying-sociocultural-barriers-and-facilitating-factors-nutrition-related-behavior>

Impact evaluation of the DFID Programme to accelerate improved nutrition for the extreme poor in Bangladesh: Final Report

URL: <http://refinenutrition.org/content/impact-evaluation-dfid-programme-accelerate-improved-nutrition-extreme-poor-bangladesh-final>

Impact of daily versus weekly supply of locally produced ready-to-use food on growth of moderately wasted children on Nias Island, Indonesia

URL: <http://refinenutrition.org/content/impact-daily-versus-weekly-supply-locally-produced-ready-use-food-growth-moderately-wasted>

Impact of facility based dietary management on severe acute malnutrition in children aged 6 to 60 months admitted in nutritional rehabilitation centre of KLESs Dr. Prabhakar Kore Hospital of Belagavi City - A longitudinal study

URL: <http://refinenutrition.org/content/impact-facility-based-dietary-management-severe-acute-malnutrition-children-aged-6-60-months>

Impact of fortified blended food aid products on nutritional status of infants and young children in developing countries

URL: <http://refinenutrition.org/content/impact-fortified-blended-food-aid-products-nutritional-status-infants-and-young-children>

Impact of lipid-based nutrient supplementation (LNS) on children's diet adequacy in Western Uganda

URL: <http://refinenutrition.org/content/impact-lipid-based-nutrient-supplementation-lns-childrens-diet-adequacy-western-uganda>

Impact of lipid-based nutrient supplements and corn-soy blend on energy and nutrient intake among moderately underweight 8-18-month-old children participating in a clinical trial.

URL: <http://refinenutrition.org/content/impact-lipid-based-nutrient-supplements-and-corn-soy-blend-energy-and-nutrient-intake-among>

Impact of therapeutic food compared to oral nutritional supplements on nutritional outcomes in mildly underweight healthy children in a low-medium income society

URL: <http://refinenutrition.org/content/impact-therapeutic-food-compared-oral-nutritional-supplements-nutritional-outcomes-mildly>

Impact study report: Comparison of PM2A and MCHN approaches in SOUHARDO II

URL: <http://refinenutrition.org/content/impact-study-report-comparison-pm2a-and-mchn-approaches-souhar-do-ii>

Impaired bile acid homeostasis in children with severe acute malnutrition

URL: <http://refinenutrition.org/content/impaired-bile-acid-homeostasis-children-severe-acute-malnutrition>

Improve nutrition, food safety, and food security for Chinas most vulnerable women and children Joint final evaluation

URL: <http://refinenutrition.org/content/improve-nutrition-food-safety-and-food-security-chinas-most-vulnerable-women-and-children->

In-home fortification with 2.5mg iron as NaFeEDTA does not reduce anaemia but increases weight gain: a randomised controlled trial in Kenyan infants

URL: <http://refinenutrition.org/content/home-fortification-25mg-iron-nafeedta-does-not-reduce-anaemia-increases-weight-gain>

Including whey protein and whey permeate in ready-to-use supplementary food improves recovery rates in children with moderate acute malnutrition: a randomized, double-blind clinical trial

URL: <http://refinenutrition.org/content/including-whey-protein-and-whey-permeate-ready-use-supplementary-food-improves-recovery>

Influence of prenatal and postnatal growth on intellectual functioning in school-aged children

URL: <http://refinenutrition.org/content/influence-prenatal-and-postnatal-growth-intellectual-functioning-school-aged-children>

Intervention study shows suboptimal growth among children receiving a food supplement for five months in a slum in Bangladesh

URL: <http://refinenutrition.org/content/intervention-study-shows-suboptimal-growth-among-children-receiving-food-supplement-five>

Interventions to improve intake of complementary foods by infants 6 to 12 months of age in developing countries: impact on growth and on the prevalence of malnutrition and potential contribution to child survival

URL: <http://refinenutrition.org/content/interventions-improve-intake-complementary-foods-infants-6-12-months-age-developing>

Intra-household use and acceptability of Ready-to-Use-Supplementary-Foods distributed in Niger between July and December 2010

URL: <http://refinenutrition.org/content/intra-household-use-and-acceptability-ready-use-supplementary-foods-distributed-niger>

Investigation of food acceptability and feeding practices for lipid nutrient supplements and blended flours used to treat moderate malnutrition

URL: <http://refinenutrition.org/content/investigation-food-acceptability-and-feeding-practices-lipid-nutrient-supplements-and>

Iron bioavailability from a lipid-based complementary food fortificant mixed with millet porridge can be optimized by adding phytase and ascorbic acid but not by using a mixture of ferrous sulfate and sodium iron EDTA

URL: <http://refinenutrition.org/content/iron-bioavailability-lipid-based-complementary-food-fortificant-mixed-millet-porridge-can-be>

Iron bioavailability in 8-24-month-old Thai children from a micronutrient-fortified quick-cooking rice containing ferric ammonium citrate or a mixture of ferrous sulphate and ferric sodium ethylenediaminetetraacetic acid

URL: <http://refinenutrition.org/content/iron-bioavailability-8-24-month-old-thai-children-micronutrient-fortified-quick-cooking-rice>

Iron fortification adversely affects the gut microbiome, increases pathogen abundance and induces intestinal inflammation in Kenyan infants

URL: <http://refinenutrition.org/content/iron-fortification-adversely-affects-gut-microbiome-increases-pathogen-abundance-and-induces>

Is the inclusion of animal source foods in fortified blended food justified?

URL: <http://refinenutrition.org/content/inclusion-animal-source-foods-fortified-blended-food-justified>

Linear growth increased in young children in an urban slum of Haiti: a randomized controlled trial of a lipid-based nutrient supplement

URL: <http://refinenutrition.org/content/linear-growth-increased-young-children-urban-slum-haiti-randomized-controlled-trial-lipid>

Lipid-based nutrient supplement increases the birth size of infants of primiparous women in Ghana

URL: <http://refinenutrition.org/content/lipid-based-nutrient-supplement-increases-birth-size-infants-primiparous-women-ghana>

Lipid-based nutrient supplementation in the first 1000 d improves child growth in Bangladesh: a cluster-randomized effectiveness trial

URL: <http://refinenutrition.org/content/lipid-based-nutrient-supplementation-first-1000-d-improves-child-growth-bangladesh-cluster>

Lipid-based nutrient supplements are feasible as a breastmilk replacement for HIV-exposed infants from 24 to 48 weeks of age

URL: <http://refinenutrition.org/content/lipid-based-nutrient-supplements-are-feasible-breastmilk-replacement-hiv-exposed-infants-24>

Lipid-based nutrient supplements for pregnant women reduce newborn stunting in a cluster-randomized controlled effectiveness trial in Bangladesh

URL: <http://refinenutrition.org/content/lipid-based-nutrient-supplements-pregnant-women-reduce-newborn-stunting-cluster-randomized>

Lipid-based nutrient supplements increase energy and macronutrient intakes from complementary food among Malawian infants

URL: <http://refinenutrition.org/content/lipid-based-nutrient-supplements-increase-energy-and-macronutrient-intakes-complementary>

Lipid-based nutrient supplements: how can they combat child malnutrition?

URL: <http://refinenutrition.org/content/lipid-based-nutrient-supplements-how-can-they-combat-child-malnutrition>

Local production of ready-to-use therapeutic food for the treatment of severe acute malnutrition

URL: <http://refinenutrition.org/content/local-production-ready-use-therapeutic-food-treatment-severe-acute-malnutrition>

Local versus offshore production of ready-to-use therapeutic foods and small quantity lipid-based nutrient supplements

URL: <http://refinenutrition.org/content/local-versus-offshore-production-ready-use-therapeutic-foods-and-small-quantity-lipid-based>

Locally made ready to use therapeutic food for treatment of malnutrition a randomized controlled trial

URL: <http://refinenutrition.org/content/locally-made-ready-use-therapeutic-food-treatment-malnutrition-randomized-controlled-trial>

Locally produced cereal/nut/legume-based biscuits versus peanut/milk-based spread for treatment of moderately to mildly wasted children in daily programmes on Nias Island, Indonesia: an issue of acceptance and compliance?

URL: <http://refinenutrition.org/content/locally-produced-cereal-nut-legume-based-biscuits-versus-peanut-milk-based-spread-treatment>

Longitudinal growth and health outcomes in nutritionally at-risk children who received long-term nutritional intervention

URL: <http://refinenutrition.org/content/longitudinal-growth-and-health-outcomes-nutritionally-risk-children-who-received-long-term>

Low-cost, ready-to-use therapeutic foods can be designed using locally available commodities with the aid of linear programming

URL: <http://refinenutrition.org/content/low-cost-ready-use-therapeutic-foods-can-be-designed-using-locally-available-commodities-aid>

Lowdose RUTF protocol and improved service delivery lead to good programme outcomes in the treatment of uncomplicated SAM: a programme report from Myanmar

URL: <http://refinenutrition.org/content/lowdose-rutf-protocol-and-improved-service-delivery-lead-good-programme-outcomes-treatment>

Malawian mothers consider lipid-based nutrient supplements acceptable for children throughout a 1-year intervention, but deviation from user recommendations is common

URL: <http://refinenutrition.org/content/malawian-mothers-consider-lipid-based-nutrient-supplements-acceptable-children-throughout-1>

Malawian mothers' attitudes towards the use of two supplementary foods for moderately malnourished children

URL: <http://refinenutrition.org/content/malawian-mothers-attitudes-towards-use-two-supplementary-foods-moderately-malnourished>

Malian children with moderate acute malnutrition who are treated with lipid-based dietary supplements have greater weight gains and recovery rates than those treated with locally produced cereal-legume products: a community-based, cluster-randomized

URL: <http://refinenutrition.org/content/malian-children-moderate-acute-malnutrition-who-are-treated-lipid-based-dietary-supplements>

Managing acute malnutrition: A review of the evidence and country experiences in South Asia and a recommended approach for Bangladesh

URL: <http://refinenutrition.org/content/managing-acute-malnutrition-review-evidence-and-country-experiences-south-asia-and>

Meat as complementary food for older breastfed infants and toddlers: a randomized, controlled trial in rural China

URL: <http://refinenutrition.org/content/meat-complementary-food-older-breastfed-infants-and-toddlers-randomized-controlled-trial>

Micronutrient supplementation, dietary intervention and resulting body weight gain of severe acute malnourished children: a pilot project study of OJUS Medical Institute

URL: <http://refinenutrition.org/content/micronutrient-supplementation-dietary-intervention-and-resulting-body-weight-gain-severe>

Micronutrients in the treatment of stunting and moderate malnutrition

URL: <http://refinenutrition.org/content/micronutrients-treatment-stunting-and-moderate-malnutrition>

Milk and Dairy Products in Human Nutrition

URL: <http://refinenutrition.org/content/milk-and-dairy-products-human-nutrition>

Milk and Dairy Products in Human Nutrition

URL: <http://refinenutrition.org/content/milk-and-dairy-products-human-nutrition>

Most children who took part in a comprehensive malnutrition programme in Madagascar reached and maintained the recovery threshold

URL: <http://refinenutrition.org/content/most-children-who-took-part-comprehensive-malnutrition-programme-madagascar-reached-and>

Multi-micronutrient fortified rice improved vitamin A status of Cambodian school children

URL: <http://refinenutrition.org/content/multi-micronutrient-fortified-rice-improved-vitamin-status-cambodian-school-children>

Multiple micronutrient powders for home (point of use) fortification of foods in pregnant women: a systematic review

URL: <http://refinenutrition.org/content/multiple-micronutrient-powders-home-point-use-fortification-foods-pregnant-women-systematic>

Nutrition in emergencies: Do we know what works?

URL: <http://refinenutrition.org/content/nutrition-emergencies-do-we-know-what-works>

Nutrition interventions for children aged less than 5 years following natural disasters: a systematic review

URL: <http://refinenutrition.org/content/nutrition-interventions-children-aged-less-5-years-following-natural-disasters-systematic>

Nutrition rehabilitation of undernourished children utilizing Spiruline and Misola

URL: <http://refinenutrition.org/content/nutrition-rehabilitation-undernourished-children-utilizing-spiruline-and-misola>

Optimisation of a cheap and residential smallscale production of edible crickets with local byproducts as an alternative proteinrich human food source in Ratanakiri Province, Cambodia

URL: <http://refinenutrition.org/content/optimisation-cheap-and-residential-smallscale-production-edible-crickets-local-products>

Optimizing ready-to-use therapeutic foods for protein quality, cost, and acceptability

URL: <http://refinenutrition.org/content/optimizing-ready-use-therapeutic-foods-protein-quality-cost-and-acceptability>

Optimizing ready-to-use therapeutic foods for protein quality, cost, and acceptability

URL: <http://refinenutrition.org/content/optimizing-ready-use-therapeutic-foods-protein-quality-cost-and-acceptability>

Outpatient treatment of severe acute malnutrition: response to treatment with a reduced schedule of therapeutic food distribution

URL: <http://refinenutrition.org/content/outpatient-treatment-severe-acute-malnutrition-response-treatment-reduced-schedule>

Patterns and determinants of small-quantity LNS utilization in rural Malawi and Mozambique: considerations for interventions with specialized nutritious foods

URL: <http://refinenutrition.org/content/patterns-and-determinants-small-quantity-lns-utilization-rural-malawi-and-mozambique>

Peanut-based ready-to-use therapeutic food: acceptability among malnourished children and community workers in Bangladesh

URL: <http://refinenutrition.org/content/peanut-based-ready-use-therapeutic-food-acceptability-among-malnourished-children-and>

Peanut-based ready-to-use therapeutic food: How acceptable and tolerated is it among malnourished pregnant and lactating women in Bangladesh?

URL: <http://refinenutrition.org/content/peanut-based-ready-use-therapeutic-food-how-acceptable-and-tolerated-it-among-malnourished>

Perceptions of usage and unintended consequences of provision of ready-to-use therapeutic food for management of severe acute child malnutrition. A qualitative study in Southern Ethiopia

URL: <http://refinenutrition.org/content/perceptions-usage-and-unintended-consequences-provision-ready-use-therapeutic-food>

Performance of UNHCR nutrition programs in post-emergency refugee camps

URL: <http://refinenutrition.org/content/performance-unhcr-nutrition-programs-post-emergency-refugee-camps>

Persistent gut microbiota immaturity in malnourished Bangladeshi children

URL: <http://refinenutrition.org/content/persistent-gut-microbiota-immaturity-malnourished-bangladeshi-children>

Persistent gut microbiota immaturity in malnourished Bangladeshi children

URL: <http://refinenutrition.org/content/persistent-gut-microbiota-immaturity-malnourished-bangladeshi-children>

Postintervention growth of Malawian children who received 12-mo dietary complementation with a lipid-based nutrient supplement or maize-soy flour

URL: <http://refinenutrition.org/content/postintervention-growth-malawian-children-who-received-12-mo-dietary-complementation-lipid>

Preferred delivery method and acceptability of Wheat-Soy Blend (WSB++) as a daily complementary food supplement in northwest Bangladesh

URL: <http://refinenutrition.org/content/preferred-delivery-method-and-acceptability-wheat-soy-blend-wsb-daily-complementary-food>

Preventative lipid-based nutrient supplements (LNS) and young child feeding practices: findings from qualitative research in Haiti

URL: <http://refinenutrition.org/content/preventative-lipid-based-nutrient-supplements-lns-and-young-child-feeding-practices-findings>

Preventing acute malnutrition among young children in crises: A prospective intervention study in Niger

URL: <http://refinenutrition.org/content/preventing-acute-malnutrition-among-young-children-crises-prospective-intervention-study-0>

Preventing acute malnutrition in young children: Improving the evidence for current and future practice

URL: <http://refinenutrition.org/content/preventing-acute-malnutrition-young-children-improving-evidence-current-and-future-practice>

Prevention of acute malnutrition: distribution of special nutritious foods and cash, and addressing underlying causes--what to recommend when, where, for whom, and how

URL: <http://refinenutrition.org/content/prevention-acute-malnutrition-distribution-special-nutritious-foods-and-cash-and-addressing>

Prevention of acute malnutrition: distribution of special nutritious foods and cash, and addressing underlying causes--what to recommend when, where, for whom, and how

URL: <http://refinenutrition.org/content/prevention-acute-malnutrition-distribution-special-nutritious-foods-and-cash-and-addressing>

Preventive effects of long-term supplementation with 2 nutritious food supplements in young children in Niger

URL: <http://refinenutrition.org/content/preventive-effects-long-term-supplementation-2-nutritious-food-supplements-young-children>

Probiotics and prebiotics for severe acute malnutrition (PRONUT study): a double-blind efficacy randomised controlled trial in Malawi

URL: <http://refinenutrition.org/content/probiotics-and-prebiotics-severe-acute-malnutrition-pronut-study-double-blind-efficacy>

Program changes are effective and cost-effective in increasing the amount of oil used in preparing corn soy blend porridge for treatment of moderate acute malnutrition in Malawi

URL: <http://refinenutrition.org/content/program-changes-are-effective-and-cost-effective-increasing-amount-oil-used-preparing-corn>

Protein quality and growth in malnourished children

URL: <http://refinenutrition.org/content/protein-quality-and-growth-malnourished-children>

Providing lipid-based nutrient supplements does not affect developmental milestones among Malawian children

URL: <http://refinenutrition.org/content/providing-lipid-based-nutrient-supplements-does-not-affect-developmental-milestones-among>

Provision of 1040 g/d lipid-based nutrient supplements from 6 to 18 months of age does not prevent linear growth faltering in Malawi

URL: <http://refinenutrition.org/content/provision-1040-gd-lipid-based-nutrient-supplements-6-18-months-age-does-not-prevent-linear>

Provision of lipid-based nutrient supplements to Honduran children increases their dietary macro- and micronutrient intake without displacing other foods.

URL: <http://refinenutrition.org/content/provision-lipid-based-nutrient-supplements-honduran-children-increases-their-dietary-macro-0>

Quantifying and exploiting the age dependence in the effect of supplementary food for child undernutrition

URL: <http://refinenutrition.org/content/quantifying-and-exploiting-age-dependence-effect-supplementary-food-child-undernutrition>

Randomized controlled trial of the effectiveness of a soybean-maize-sorghum-based ready-to-use complementary food paste on infant growth in South Kivu, Democratic Republic of Congo

URL: <http://refinenutrition.org/content/randomized-controlled-trial-effectiveness-soybean-maize-sorghum-based-ready-use>

Randomized controlled trial of the impact of treating moderately malnourished women in pregnancy

URL: <http://refinenutrition.org/content/randomized-controlled-trial-impact-treating-moderately-malnourished-women-pregnancy>

Ready-to-use food-allocation policy to reduce the effects of childhood undernutrition in developing countries

URL: <http://refinenutrition.org/content/ready-use-food-allocation-policy-reduce-effects-childhood-undernutrition-developing>

Ready-to-use foods for management of moderate acute malnutrition: Considerations for scaling up production and use in programs

URL: <http://refinenutrition.org/content/ready-use-foods-management-moderate-acute-malnutrition-considerations-scaling-production-and>

Ready-to-use supplementary food increases fat mass and BMI in Haitian school-aged children

URL: <http://refinenutrition.org/content/ready-use-supplementary-food-increases-fat-mass-and-bmi-haitian-school-aged-children>

Ready-to-use supplementary food versus corn soya blend with oil premix to treat moderate acute child malnutrition: a community-based cluster randomized trial

URL: <http://refinenutrition.org/content/ready-use-supplementary-food-versus-corn-soya-blend-oil-premix-treat-moderate-acute-child>

Ready-to-use supplementary foods and ready-to-use therapeutic foods developing product standards

URL: <http://refinenutrition.org/content/ready-use-supplementary-foods-and-ready-use-therapeutic-foods-developing-product-standards>

Ready-to-use therapeutic food for home-based treatment of severe acute malnutrition in children from six months to five years of age

URL: <http://refinenutrition.org/content/ready-use-therapeutic-food-home-based-treatment-severe-acute-malnutrition-children-six>

Ready-to-use therapeutic food for home-based treatment of severe acute malnutrition in children from six months to five years of age

URL: <http://refinenutrition.org/content/ready-use-therapeutic-food-home-based-treatment-severe-acute-malnutrition-children-six>

Ready-to-use therapeutic food with elevated n-3 polyunsaturated fatty acid content, with or without fish oil, to treat severe acute malnutrition: a randomized controlled trial

URL: <http://refinenutrition.org/content/ready-use-therapeutic-food-elevated-n-3-polyunsaturated-fatty-acid-content-or-without-fish>

Reasons for raising the maximum acceptable daily intake of EDTA and the benefits for iron fortification of foods for children 6-24 months of age

URL: <http://refinenutrition.org/content/reasons-raising-maximum-acceptable-daily-intake-edta-and-benefits-iron-fortification-foods>

Recovery rate and determinants in treatment of children with severe acute malnutrition using outpatient therapeutic feeding program in Kamba District, Ethiopia

URL: <http://refinenutrition.org/content/recovery-rate-and-determinants-treatment-children-severe-acute-malnutrition-using-outpatient>

Recovery rate of children with moderate acute malnutrition treated with ready-to-use supplementary food (RUSF) or improved cornsoya blend (CSB+): a randomized controlled trial

URL: <http://refinenutrition.org/content/recovery-rate-children-moderate-acute-malnutrition-treated-ready-use-supplementary-food-rusf>

Recovery rate of children with moderate acute malnutrition treated with ready-to-use supplementary food (RUSF) or improved cornsoya blend (CSB+): a randomized controlled trial

URL: <http://refinenutrition.org/content/recovery-rate-children-moderate-acute-malnutrition-treated-ready-use-supplementary-food-rusf>

Recovery rate of children with moderate acute malnutrition treated with ready-to-use supplementary food (RUSF) or improved cornsoya blend (CSB+): a randomized controlled trial

URL: <http://refinenutrition.org/content/recovery-rate-children-moderate-acute-malnutrition-treated-ready-use-supplementary-food-rusf>

Reducing stunting by improving maternal, infant and young child nutrition in regions such as South Asia: evidence, challenges and opportunities

URL: <http://refinenutrition.org/content/reducing-stunting-improving-maternal-infant-and-young-child-nutrition-regions-such-south>

Research gaps in the use of dairy ingredients in food aid products

URL: <http://refinenutrition.org/content/research-gaps-use-dairy-ingredients-food-aid-products>

Research priorities for improving infant and young child feeding in humanitarian emergencies

URL: <http://refinenutrition.org/content/research-priorities-improving-infant-and-young-child-feeding-humanitarian-emergencies>

Responsive feeding and child interest in food vary when rural Malawian children are fed lipidbased nutrient supplements or local complementary food

URL: <http://refinenutrition.org/content/responsive-feeding-and-child-interest-food-vary-when-rural-malawian-children-are-fed-lipid>

Review Protocol: Food fortification with multiple micronutrients: impact on health

URL: <http://refinenutrition.org/content/review-protocol-food-fortification-multiple-micronutrients-impact-health>

Risk factors for death in children during inpatient treatment of severe acute malnutrition: a prospective cohort study

URL: <http://refinenutrition.org/content/risk-factors-death-children-during-inpatient-treatment-severe-acute-malnutrition-prospective>

Routine amoxicillin for uncomplicated severe acute malnutrition in children

URL: <http://refinenutrition.org/content/routine-amoxicillin-uncomplicated-severe-acute-malnutrition-children>

Screening for anti-nutritional compounds in complementary foods and food aid products for infants and young children

URL: <http://refinenutrition.org/content/screening-anti-nutritional-compounds-complementary-foods-and-food-aid-products-infants-and>

Seasonal effect and long-term nutritional status following exit from a Community-Based Management of Severe Acute Malnutrition program in Bihar, India

URL: <http://refinenutrition.org/content/seasonal-effect-and-long-term-nutritional-status-following-exit-community-based-management>

Seasonal effect and long-term nutritional status following exit from a Community-Based Management of Severe Acute Malnutrition program in Bihar, India

URL: <http://refinenutrition.org/content/seasonal-effect-and-long-term-nutritional-status-following-exit-community-based-management>

Seasonality modifies the effect of a lipid-based nutrient supplement for pregnant rural women on birth length

URL: <http://refinenutrition.org/content/seasonality-modifies-effect-lipid-based-nutrient-supplement-pregnant-rural-women-birth>

Severe acute malnutrition

URL: <http://refinenutrition.org/content/severe-acute-malnutrition>

Severe and moderate acute malnutrition can be successfully managed with an integrated protocol in Sierra Leone

URL: <http://refinenutrition.org/content/severe-and-moderate-acute-malnutrition-can-be-successfully-managed-integrated-protocol>

Short children with a low midupper arm circumference respond to food supplementation: an observational study from Burkina Faso

URL: <http://refinenutrition.org/content/short-children-low-midupper-arm-circumference-respond-food-supplementation-observational>

Small-quantity lipid-based nutrient supplements, regardless of their zinc content, increase growth and reduce the prevalence of stunting and wasting in young burkinabe children: a cluster-randomized trial

URL: <http://refinenutrition.org/content/small-quantity-lipid-based-nutrient-supplements-regardless-their-zinc-content-increase>

Small-quantity, lipid-based nutrient supplements provided to women during pregnancy and 6 mo postpartum and to their infants from 6 mo of age increase the mean attained length of 18-mo-old children in semi-urban Ghana: a randomized controlled trial

URL: <http://refinenutrition.org/content/small-quantity-lipid-based-nutrient-supplements-provided-women-during-pregnancy-and-6-mo>

Socio-anthropological investigation related to the acceptability of Plumpy'nut in Cambodia

URL: <http://refinenutrition.org/content/socio-anthropological-investigation-related-acceptability-plumpynut-cambodia>

Specially formulated foods for treating children with moderate acute malnutrition in low-and middle-income countries

URL: <http://refinenutrition.org/content/specially-formulated-foods-treating-children-moderate-acute-malnutrition-low-and-middle>

Strengthening and evaluating the Preventing Malnutrition in Children under 2 Years of Age approach Burundi Follow-up Report: Children 023 Months

URL: <http://refinenutrition.org/content/strengthening-and-evaluating-preventing-malnutrition-children-under-2-years-age-approach->

Studying the effectiveness of mixed models of community based and institutional based interventions for management of severe acute malnourished children

URL: <http://refinenutrition.org/content/studying-effectiveness-mixed-models-community-based-and-institutional-based-interventions>

Supplementary feeding with fortified spread among moderately underweight 6-18-month-old rural Malawian children

URL: <http://refinenutrition.org/content/supplementary-feeding-fortified-spread-among-moderately-underweight-6-18-month-old-rural>

Supplementary feeding with fortified spread among moderately underweight 6-18-month-old rural Malawian children

URL: <http://refinenutrition.org/content/supplementary-feeding-fortified-spread-among-moderately-underweight-6-18-month-old-rural>

Supplementary feeding with fortified spreads results in higher recovery rates than with a corn/soy blend in moderately wasted children

URL: <http://refinenutrition.org/content/supplementary-feeding-fortified-spreads-results-higher-recovery-rates-cornsoy-blend>

Supplementary feeding with locally-produced ready-to-use food (RUF) for mildly wasted children on Nias island, Indonesia: Comparison of daily and weekly program outcomes

URL: <http://refinenutrition.org/content/supplementary-feeding-locally-produced-ready-use-food-ruf-mildly-wasted-children-nias-island>

Supplementation of maternal diets during pregnancy and for 6 months postpartum and infant diets thereafter with small-quantity lipid-based nutrient supplements does not promote child growth by 18 months of age in rural Malawi: A randomized controlled trial

URL: <http://refinenutrition.org/content/supplementation-maternal-diets-during-pregnancy-and-6-months-postpartum-and-infant-diets>

Systematic review of the efficacy and effectiveness of complementary feeding interventions in developing countries

URL: <http://refinenutrition.org/content/systematic-review-efficacy-and-effectiveness-complementary-feeding-interventions-developing>

The effect of adding ready-to-use supplementary food to a general food distribution on child nutritional status and morbidity: a cluster-randomized controlled trial

URL: <http://refinenutrition.org/content/effect-adding-ready-use-supplementary-food-general-food-distribution-child-nutritional>

The effect of balanced protein energy supplementation in undernourished pregnant women and child physical growth in low- and middle-income countries: a systematic review and meta-analysis

URL: <http://refinenutrition.org/content/effect-balanced-protein-energy-supplementation-undernourished-pregnant-women-and-child>

The effect of low dose iron and zinc intake on child micronutrient status and development during the first 1000 days of life: A systematic review and meta-analysis

URL: <http://refinenutrition.org/content/effect-low-dose-iron-and-zinc-intake-child-micronutrient-status-and-development-during-first>

The effect of supplementation with locally available foods on stunting: A review of theory and evidence

URL: <http://refinenutrition.org/content/effect-supplementation-locally-available-foods-stunting-review-theory-and-evidence>

The effect of supplementation with locally available foods on stunting: A review of theory and evidence

URL: <http://refinenutrition.org/content/effect-supplementation-locally-available-foods-stunting-review-theory-and-evidence>

The effectiveness of interventions to treat severe acute malnutrition in young children: a systematic review

URL: <http://refinenutrition.org/content/effectiveness-interventions-treat-severe-acute-malnutrition-young-children-systematic-review>

The effects of iron fortification on the gut microbiota in African children: a randomized controlled trial in Cote d'Ivoire

URL: <http://refinenutrition.org/content/effects-iron-fortification-gut-microbiota-african-children-randomized-controlled-trial-cote>

The impact and effectiveness of emergency nutrition and nutrition related interventions: a review of published evidence 2004-2010

URL: <http://refinenutrition.org/content/impact-and-effectiveness-emergency-nutrition-and-nutrition-related-interventions-review>

The impact of dietary supplementation with lipid-based nutrient supplements on maternal health and pregnancy outcomes in rural Malawi

URL: <http://refinenutrition.org/content/impact-dietary-supplementation-lipid-based-nutrient-supplements-maternal-health-and>

The impact of integrated infant and young child feeding and micronutrient powder intervention on feeding practices and anemia in children aged 6-23 months in Madagascar

URL: <http://refinenutrition.org/content/impact-integrated-infant-and-young-child-feeding-and-micronutrient-powder-intervention>

The impact of lipid-based nutrient supplement provision to pregnant women on newborn size in rural Malawi: a randomized controlled trial

URL: <http://refinenutrition.org/content/impact-lipid-based-nutrient-supplement-provision-pregnant-women-newborn-size-rural-malawi>

The impact of Ready-to-Use Supplementary Food (RUSF) in targeted supplementation of children with moderate acute malnutrition (MAM) in South Africa

URL: <http://refinenutrition.org/content/impact-ready-use-supplementary-food-rusf-targeted-supplementation-children-moderate-acute>

The MAM'Out project: a randomized controlled trial to assess multiannual and seasonal cash transfers for the prevention of acute malnutrition in children

URL: <http://refinenutrition.org/content/mamout-project-randomized-controlled-trial-assess-multiannual-and-seasonal-cash-transfers>

The price of oil: Behavior change communication and increased oil ration improve mothers' uses of vegetable oil in preparing food aid rations for children in Malawi

URL: <http://refinenutrition.org/content/price-oil-behavior-change-communication-and-increased-oil-ration-improve-mothers-uses>

The REFANI Pakistan study-a cluster randomised controlled trial of the effectiveness and cost-effectiveness of cash-based transfer programmes on child nutrition status: study protocol

URL: <http://refinenutrition.org/content/refani-pakistan-study-cluster-randomised-controlled-trial-effectiveness-and-cost>

The role of dairy in comparative effectiveness and cost of treatment of children with moderate acute malnutrition: A narrative review

URL: <http://refinenutrition.org/content/role-dairy-comparative-effectiveness-and-cost-treatment-children-moderate-acute-malnutrition>

The role of dairy in the comparative effectiveness and cost of fortified blended foods versus ready-to-use foods in treatment of children with moderate acute malnutrition

URL: <http://refinenutrition.org/content/role-dairy-comparative-effectiveness-and-cost-fortified-blended-foods-versus-ready-use-foods>

The use of linear programming to determine whether a formulated complementary food product can ensure adequate nutrients for 6- to 11-month-old Cambodian infants

URL: <http://refinenutrition.org/content/use-linear-programming-determine-whether-formulated-complementary-food-product-can-ensure>

The use of nutrient-optimizing/cost-minimizing software to develop ready-to-use therapeutic foods for malnourished pregnant women in Mali

URL: <http://refinenutrition.org/content/use-nutrient-optimizingcost-minimizing-software-develop-ready-use-therapeutic-foods>

Treating moderate acute malnutrition in first-line health services: an effectiveness cluster-randomized trial in Burkina Faso

URL: <http://refinenutrition.org/content/treating-moderate-acute-malnutrition-first-line-health-services-effectiveness-cluster>

Treatment of moderate acute malnutrition with ready-to-use supplementary food results in higher overall recovery rates compared with a corn-soya blend in children in southern Ethiopia: an operations research trial

URL: <http://refinenutrition.org/content/treatment-moderate-acute-malnutrition-ready-use-supplementary-food-results-higher-overall>

Treatment of severe and moderate acute malnutrition in low-and middle-income settings: a systematic review, meta-analysis and Delphi process

URL: <http://refinenutrition.org/content/treatment-severe-and-moderate-acute-malnutrition-low-and-middle-income-settings-systematic>

Treatment of severe and moderate acute malnutrition in low-and middle-income settings: a systematic review, meta-analysis and Delphi process

URL: <http://refinenutrition.org/content/treatment-severe-and-moderate-acute-malnutrition-low-and-middle-income-settings-systematic>

Tubaramure, a food-assisted integrated health and nutrition program in Burundi, increases maternal and child hemoglobin concentrations and reduces anemia: A theory-based cluster-randomized controlled intervention trial

URL: <http://refinenutrition.org/content/tubaramure-food-assisted-integrated-health-and-nutrition-program-burundi-increases-maternal>

Unconditional cash transfers do not prevent children's undernutrition in the Moderate Acute Malnutrition Out (MAM'Out) cluster-randomized controlled trial in rural Burkina Faso

URL: <http://refinenutrition.org/content/unconditional-cash-transfers-do-not-prevent-childrens-undernutrition-moderate-acute>

Whole cowpea meal fortified with NaFeEDTA reduces iron deficiency among Ghanaian school children in a malaria endemic area

URL: <http://refinenutrition.org/content/whole-cowpea-meal-fortified-nafeedta-reduces-iron-deficiency-among-ghanaian-school-children>

Willingness to pay for lipidbased nutrient supplements for young children in four urban sites of Ethiopia

URL: <http://refinenutrition.org/content/willingness-pay-lipidbased-nutrient-supplements-young-children-four-urban-sites-ethiopia>

This website is made possible through support of the USAID's Bureau for Democracy, Conflict and Humanitarian Assistance in the Office of Food For Peace. The contents are the responsibility of Tufts University and do not necessarily reflect the views of the U.S. Agency for International Development.